

**EPB785Hu61 100µg**

**Eukaryotic Neprilysin (CD10)**

**Organism Species: *Homo sapiens* (Human)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

12th Edition (Revised in Aug, 2016)

**[ PROPERTIES ]**

**Source:** Eukaryotic expression

**Host:** 293F cell

**Residues:** Tyr52~Trp750

**Tags:** N-terminal His Tag

**Subcellular Location:** Membrane

**Purity:** > 95%

**Traits:** Freeze-dried powder

**Buffer formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% SKL, 5% Trehalose.

**Original Concentration:** 250µg/mL

**Applications:** Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 5.2

**Predicted Molecular Mass:** 81.6kDa

**Accurate Molecular Mass:** 100kDa as determined by SDS-PAGE reducing conditions.

**Phenomenon explanation:**

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affect the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

**[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

**[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


**[ SEQUENCE ]**

```

YDDGICKSS DCIKSAARLI QNMDATTEPC TDDFKYACGG WLKRVIPET
SSRYGNFDIL RDELEVVLKD VLQEPKTEDI VAVQKAKALY RSCINESAID
SRGGEPLLLK LPDIYGWVPA TENWEQKYGA SWTAEKAIQA LNSKYGKKVL
INLFGTDDK NSVNHVIHID QPRLGLPSRD YYECTGIYKE ACTAYVDFMI
SVARLIRQEE RLPIDENQLA LEMNKVMELE KEIANATAKP EDRNDPMLLY
NKMTLAQIQN NFSLEINGKP FSWLNFTNEI MSTVNISITN EEDVVVYAPE
YLTCLKPILT KYSARDLQNL MSWRFIMDLV SLSRPTYKES RNAFRKALYG
TTSETATWRR CANYVNGNME NAVGRLYVEA AFAGESKHVV EDLIAQIREV
FIQTLDDLTW MDAETKKRAE EKALAIKERI GYPDDIVSND NKLNNEYLEL
NYKEDEYFEN IIQNLKFSQS KQLKKLREKV DKDEWISGAA VVNAFYSSGR
NQIVFPAGIL QPPFFSAQQS NSLNYYGIGM VIGHEITHGF DDNGRNFNKD
GDLVDWWTQQ SASNFKEQSQ CMVYQYGNFS WDLAGGQHLN GINTLGENIA
DNGGLGQAYR AYQNYIKKNG EEKLLPGLDL NHKQLFFLNF AQVWCGTYRP
EYAVNSIKTD VHSPGNFRII GTLQNSAEFS EAFHCRKNSY MNPEKKCRVW
    
```

**[ IDENTIFICATION ]**

CGATCTAGCATGATGATGATTTGGAGTCACTGACTGCAAAATCGCTGCTGCTGACTGATCGAAGGATGCGATGCGCCACTGAGCCTTGTACGACTTTTTGAAATATGCTTGGGAGGCTGTTGAAAGTAAATGTCATTCGCGAGICGAGCTCCGTTAGGGCAGCTTTGAGATTTTAAAGGATGAACTAGAG



**Figure . Gene Sequencing (extract)**




Figure. SDS-PAGE

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.