

**APJ516Mu01 100µg**  
**Active Lecithin Cholesterol Acyltransferase (LCAT)**  
**Organism Species: *Mus musculus* (Mouse)**  
***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

---

1st Edition (Apr, 2016)

## **[ PROPERTIES ]**

**Source:** Prokaryotic expression.

**Host:** *E. coli*

**Residues:** Glu41~His210

**Tags:** N-terminal His-tag

**Purity:** >98%

**Buffer Formulation:** 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

**Applications:** Cell culture; Activity Assays; In vivo assays.

(May be suitable for use in other assays to be determined by the end user.)

**Predicted isoelectric point:** 6.3

**Predicted Molecular Mass:** 20.7kDa

**Accurate Molecular Mass:** 22kDa as determined by SDS-PAGE reducing conditions.

## **[ USAGE ]**

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCE ]**

```

                                                                    ELSNHTRPVI
LVPGCLGNRL EAKLDKPDVV NVMCYRKTED FFTIWLDFNL FLPLGVDCWI
DNTRIVYNHS SGRVSNAPGV QIRVPGFGKT ESVEYVDDNK LAGYLHTLVQ
NLVNNGYVRD ETVRAAPYDW RLAPHQQDEY YKKLGLVVEE MYAAYGKPVF
LIGHSLGCLH
```

## **[ ACTIVITY ]**

LCAT (Phosphatidylcholine-sterol acyltransferase) is an enzyme in the extracellular metabolism of plasma lipoproteins, which converts cholesterol and phosphatidylcholines (lecithins) to cholesteryl esters and lysophosphatidylcholines. It is reported that APOA1 (Apolipoprotein A-I) acts as a cofactor for the LCAT. Thus, a binding ELISA assay was conducted to detect the association of LCAT with APOA1. Briefly, recombinant rat LCAT were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100uL were then transferred to APOA1-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-LCAT pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50µL stop solution to the wells and read at 450nm immediately. The binding activity of LCAT with APOA1 was shown in Figure 1 and this effect was in a dose dependent manner.




Figure 1. The binding activity of LCAT with APOA1.

## [ IDENTIFICATION ]





Figure 2. SDS-PAGE

Sample: Active recombinant LCAT, Mouse



**Figure 3. Western Blot**

**Sample: Recombinant LCAT, Mouse;**

**Antibody: Rabbit Anti-Mouse LCAT Ab (PAJ516Mu01)**