

LAA213Po71 Biotin-Linked Polyclonal Antibody To Triggering Receptor Expressed On Myeloid Cells 1 (TREM1) Organism Species: Sus scrofa; Porcine (Pig) Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

1304 Langham Creek Dr. Suite 226, Houston, TX 77064, USA | 001-888-960-7402 | www.cloud-cloue.us | mail/g/cloud-clone.as Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail/g/cloud-clone.com

# Coud-Clone Corp.

## [PROPERTIES]

Source: Antibody labeling Host: Rabbit Purification: Antigen-specific Affinity Chromatography. Label: Biotin Original Antibody: PAA213Po01 Traits: Liquid Concentration: 200µg/mL UOM: 100µg Applications: WB; IHC; ICC.

### [IMMUNOGEN]

Immunogen: Recombinant TREM1 (GIn19~Pro238) expressed in *E.coli*. Accession No.: RPA213Po01

# [APPLICATIONS]

Western blotting: 0.5-2µg/mL Immunohistochemistry: 5-20µg/mL Immunocytochemistry: 5-20µg/mL Optimal working dilutions must be determined by end user.

# [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

# [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were



observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

1304 Langham Creek Dr, Suite 226, Huustun, TX 77084, USA | 001-888-960-7402 | www.cloud-cloue.us | mail@cloud-cloue.us Export Processing Zone, Wuhan, Hubei 430056, PRC | 0086-000-880-0687 | www.cloud-clone.com | mail@cloud-clone.com