

MAB180Hu22

**Monoclonal Antibody to Clusterin (CLU)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: 6#

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 100µl

Cross Reactivity: Porcine

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant CLU (Ser228~Glu449) expressed in E.coli

Accession No.: RPB180Hu01

#### [ APPLICATIONS ]

Western blotting: 0.01-6µg/mL;

Immunohistochemistry: 5-30µg/mL;

Immunocytochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined



by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]




Figure. Western Blot; Sample: Recombinant CLU, Human.

Western Blot; Sample: Porcine Eye	Western Blot; Sample: WERI-RB-1 cell
lysate	lysate
Primary Ab: 6µg/ml Mouse Anti-Human Primary Ab: 2µg/ml Mouse Anti-Human	

, 10	, 10
CLU Antibody	CLU Antibody
Second Ab: 0.2µg/mL HRP-Linked	Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal	Caprine Anti-Mouse IgG Polyclonal
Antibody	Antibody
(Catalog: SAA544Mu19) Selected	(Catalog: SAA544Mu19) Selected

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.