MAA726Hu22 Monoclonal Antibody to Transthyretin (TTR) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PRODUCT INFORMATION]

Corp.

Immunogen: TTR, Human Clonality: Monoclonal Clone number: C4 Host: Mouse Immunoglobulin Type: IgG1 Kappa Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 500µg/mL UOM: 200µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant TTR (Gly21~Glu147) with N-terminal His-Tag expressed in *E.coli*.

Accession No.: RPA726Hu01

[ANTIBODY SPECIFITY]

The antibody is a mouse monoclonal antibody raised against TTR. It has been selected for its ability to recognize TTR in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

Coud-Clone Corp.

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant TTR (Gly21~Glu147) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.


5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

[<u>IMAGES</u>]



Recombinant TTR, Human