

RPC282Hu01 50µg

Recombinant Adhesion Regulating Molecule 1 (ADRM1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Thr2~Asp407

Tags: N-terminal His Tag

Subcellular Location: Cytoplasm

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.01% skl, 5%Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.6

Predicted Molecular Mass: 43.4kDa

Accurate Molecular Mass: 46kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



```
TTSGALFPS LVPGSRGASN KYLVEFRAGK MSLKGTTVTP DKRKGLVYIQ
QTDDSLIHFC WKDRTSGNVE DDLIIFPDDC EFKRVPQCPS GRVYVLKFKA
GSKRLFFWMQ EPKTDQDEEH CRKVNEYLNN PPMPGALGAS GSSGHELSAL
GGEGGLQSLL GNMSHSQLMQ LIGPAGLGGL GGLGALTGPG LASLLGSSGP
PGSSSSSSR SQSAAVTPSS TTSSTRATPA PSAPAAASAT SPSPAPSSGN
GASTAASPTQ PIQLSDLQSI LATMNVPAGP AGGQQVDLAS VLTPEIMAPI
LANADVQERL LPYLPSGESL PQTADEIQNT LTSPQFQQAL GMFSAALASG
QLGPLMCQFG LPAEAVEAAN KGDVEAFAKA MQNNAKPEQK EGDTKDKKDE
EEDMSLD
```

[IDENTIFICATION]



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.