

RPE787Rb02 10µg Recombinant Annexin A1 (ANXA1) Organism Species: *Oryctolagus cuniculus (Rabbit)* Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

# Coud-Clone Corp.

### [PROPERTIES]

**Source:** Prokaryotic expression **Host:** *E.coli* 

Residues: Met1~Asn346

Tags: N-terminal His Tag

Subcellular Location: Nucleus, Cytoplasm

**Purity:** > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose .

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 6.1

Predicted Molecular Mass: 44.2kDa

Accurate Molecular Mass: 43kDa as determined by SDS-PAGE reducing conditions.

### [<u>USAGE</u>]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

### [SEQUENCE]

# Contend Clone Corp.

MAMVSEFLKQ AWFIDNEEQD YINTVKTYKG GPGSAVSPYP AFNPSSDVAA LHQAIMVKGV DEATIIDILT KRNNAQRQQI KAAYLQEKGK PLDEVLKKAL TGHLEEVVLA LLKTPAQFDA DELRAAMKGL GTDEDTLIEI LASRNNKEIR EINRVYREEL KRDLAKDIAS DTSGDFQKAL LSLAKGDRSE DFGVNEDLAD TDARALYEAG ERRKGADVNV FTTILTTRSY LHLRRVFQKY SKYSQHDMNK VLDLELKGDI EKCLTAIVQC ATCKPAYFAE KLYQAMKGAG TRHKALIRIM VSRSEVDMND IKAFYOKKYG VSLCQAILDE TKGDYEKILV ALCGGN

## [IDENTIFICATION]




Figure. SDS-PAGE

### [<u>IMPORTANT NOTE</u>]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.