EPB785Hu61 10μg

Eukaryotic Neprilysin (CD10)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Eukaryotic expression

Host: 293F cell

Residues: Tyr52~Trp750

Tags: N-terminal His Tag

Subcellular Location: Membrane

Purity: > 95%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 5% Trehalose.

Original Concentration: 200μg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.2

Predicted Molecular Mass: 81.6kDa

Accurate Molecular Mass: 100kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1.Splice variants: Alternative splicing may create different sized proteins from the same gene.

2. Relative charge: The composition of amino acids may affects the charge of the protein.

3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.

4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.

5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8ºC for one month.

Aliquot and store at -80ºC for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

YDDGICKSS DCIKSAARLI QNMDATTEPC TDFFKYACGG WLKRNVIPET
SSRYGNFDIL RDELEVVLKD VLQEPKTEDI VAVQKAKALY RSCINESAID
SRGGEPLLKL LPDIYGWPVA TENWEQKYGA SWTAEKAIAQ LNSKYGKKVL
INLFVGTDDK NSVNHVIHID QPRLGLPSRD YYECTGIYKE ACTAYVDFMI
SVARLIRQEE RLPIDENQLA LEMNKVMELE KEIANATAKP EDRNDPMLLY
NKMTLAQIQN NFSLEINGKP FSWLNFTNEI MSTVNISITN EEDVVVYAPE
YLTKLKPILT KYSARDLQNL MSWRFIMDLV SSLSRTYKES RNAFRKALYG
TTSETATWRR CANYVNGNME NAVGRLYVEA AFAGESKHVV EDLIAQIREV
FIQTLDDLTW MDAETKKRAE EKALAIKERI GYPDDIVSND NKLNNEYLEL
NYKEDEYFEN IIQNLKFSQS KQLKKLREKV DKDEWISGAA VVNAFYSSGR
NQIVFPAGIL QPPFFSAQQS NSLNYGGIGM VIGHEITHGF DDNGRNFNKD
GDLVDWWTQQ SASNFKEQSQ CMVYQYGNFS WDLAGGQHLN GINTLGENIA
DNGGLGQAYR AYQNYIKKNG EEKLLPGLDL NHKQLFFLNF AQVWCGTYRP
EYAVNSIKTD VHSPGNFRII GTLQNSAEFS EAFHCRKNSY MNPEKKCRVW

[IDENTIFICATION]



Figure . Gene Sequencing (extract)



Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.