

RPB329Hu01 10µq

Recombinant Microtubule Associated Protein 2 (MAP2)

Organism Species: Homo sapiens (Human)

Instruction manual

kDa

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES.

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Residues: Leu727~Asp985 (Accession # P11137),

with N-terminal His-Tag.

Host: E. coli

Subcellular Location: Cytoplasm, cytoskeleton.

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS, pH7.4, containing 5% sucrose, 0.01% sarcosyl.

Predicted isoelectric point: 4.7

Predicted Molecular Mass: 29.7kDa



Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note: The possible reasons that the actual band size differs from the predicted are as follows:

- 1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
- 2. Relative charge: The composition of amino acids may affects the charge of the protein.
- 3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
- 4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
- 5. Polymerization of the target protein: Dimerization, multimerization etc.



[USAGE]

Reconstitute in sterile PBS, pH7.2-pH7.4.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below. MGHHHHHHSG SEF-LASD ILTNTSGSMD EGDDYLPATT PALEKAPCFP VESKEEEQIE KVKATGEESTQAEISCESPF LAKDFYKNGT VMAPDLPEML DLAGTRSRLA SVSADAEVAR RKSVPSETVVEDSRTGLPPV TDENHVIVKT DSQLEDLGYC VFNKYTVPLP SPVQDSENLS GESGTFYEGT DDKVRRDLAT DLSLIEVKLA AAGRVKDEFS VDKEASAHIS GDKSGLSKEF DQEKKANDRLDTVLEKSEEH ADSKEHAKKT EEAGD

[RELEVANCE]

- 1. Albala J.S., et al. (1993) Gene 136:377-378.
- 2. Dammerman M., et al. (1989) J. Neurosci. Res. 24:487-495.
- 3. Kosik K.S., et al. (1988) J. Neurochem. 51:587-598.
- 4. Zamora-Leon S.P., et al. (2005) J. Biol. Chem. 280:1962-1970.