

MAB474Hu21

**Monoclonal Antibody to Nectin 2 (NECTIN2)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C6

Traits: Liquid

Concentration: 1mg/mL

**UOM:** 200µL

Cross Reactivity: N/A

Applications: WB; IHC

## [ IMMUNOGEN ]

Immunogen: Recombinant NECTIN2 (Pro76~Ala353) expressed in E.coli

Accession No.: RPB474Hu01

#### [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-40µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

# Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



Western Blot; Sample: Lane1: K562 cell

lysate; Lane2: SW480 cell lysate


Primary Ab: 2µg/ml Mouse Anti-Human

**PVRL2** Antibody

Second Ab: 0.2µg/mL HRP-Linked

Rabbit Anti-Mouse IgG Polyclonal

Antibody (Catalog: SAA544Mu19)



DAB staining on IHC-P;

Samples: Human Kidney Tissue;

Primary Ab: 40µg/ml Mouse Anti-

Human PVRL2 Antibody

Second Ab: 2µg/mL HRP-Linked

Caprine Anti-Mouse IgG Polyclonal

Antibody

(Catalog: SAA544Mu19)

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.