

MAA573Hu24

Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: J38-2

Traits: Liquid

Concentration: 1mg/mL

UOM: 100μg(100μL)

Cross Reactivity: N/A

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant PIIINP (Gln24~Pro153) expressed in E.coli

Accession No.: RPA573Hu01

[APPLICATIONS]

Western blottin:g: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no


Coud-Clone Corp.

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]




Western Blot; Sample: Lane1: Human Serum; Lane2: Human Plasma Primary Ab: 0.8µg/ml Mouse Anti-Human PIIINP Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)



DAB staining on IHC-P; Sample: 20µg/ml Mouse Anti-Human PIIINP Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

DAB staining on IHC-P; Human Placenta Tissue: Primary Ab: Sample: Human Small intestine Tissue: Primary Ab: 20µg/ml Mouse Anti-Human PIIINP Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)




DAB staining on IHC-P; Sample: Human Lymph node Tissue; Primary Ab: 20µg/ml Mouse Anti-**Human PIIINP Antibody** Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody

(Catalog: SAA544Mu19)


DAB staining on IHC-P; Sample: Human Lung Tissue; Primary Ab: 20µg/ml Mouse Anti-**Human PIIINP Antibody** Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

DAB staining on IHC-P; Sample: Human Cardiac Muscle Tissue: Primary Ab: 20µg/ml Mouse Anti-Human PIIINP Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)

Coud-Clone Corp.




DAB staining on IHC-P;
Sample: Human Uterus Tissue;
Primary Ab: 20µg/ml Mouse AntiHuman PIIINP Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P;
Sample: Human Placenta Tissue
Primary Ab: 20µg/ml Mouse AntiHuman PIIINP Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody

(Catalog: SAA544Mu19)




DAB staining on IHC-P;
Sample: Human Small intestine Tissue
Primary Ab: 20µg/ml Mouse AntiHuman PIIINP Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)



DAB staining on IHC-P;
Sample: Human Lymph node Tissue
Primary Ab: 20µg/ml Mouse AntiHuman PIIINP Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2µg/ml HRP-Linked
Caprine Anti-Mouse IgG Polyclonal

Caprine Anti-Mouse IgG Polyclonal
Antibody

(Catalog: SAA544Mu19)



Western Blot; Sample: Human Placenta
lysate
Primary Ab: 0.4µg/ml Mouse AntiHuman PIIINP Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.