

MAD227Hu21

Monoclonal Antibody to Troponin C Type 1, Slow (TNNC1)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: C5

Traits: Liquid

Concentration: 1mg/mL

UOM: 100μL

Cross Reactivity: Mouse;Porcine

Applications: WB; ICC/IF

[IMMUNOGEN]

Immunogen: Recombinant TNNC1 (Met1~Glu161) expressed in *E.coli*

Accession No.: RPD227Hu01

[APPLICATIONS]

Western blotting: 0.01-2μg/mL;

Immunocytochemistry: 5-30μg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]





Figure. Western Blot; Sample: Lane1: Human Lung Tissue; Lane2: Porcine Heart Tissue; Lane3: Porcine Skeletal Muscle Tissue; Lane4: Mouse Heart Tissue.



AF488 staining on IF;

Sample: HepG2 cell

Primary Ab: 30µg/ml Mouse Anti-Human TNNC1 Antibody

Second Ab: 2?g/ml AF488-Linked Caprine Anti-Mouse IgG Polyclonal Antibody

(Catalog: SAA544Mu11)

AF488 staining on IF;

Sample: U2OS cell

Primary Ab: 30µg/ml Mouse Anti-Human TNNC1 Antibody

Second Ab: 2?g/ml AF488-Linked Caprine Anti-Mouse IgG Polyclonal Antibody

(Catalog: SAA544Mu11)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.