

MAB070Hu23

Monoclonal Antibody to Tryptase (TPS)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2a Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: C2

Traits: Liquid

Concentration: 1mg/mL

UOM: 200μL

Cross Reactivity: Canine;Porcine

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant TPS (Gly161~Tyr262) expressed in *E.coli*

Accession No.: RPB070Hu01

[APPLICATIONS]

Western blotting: 0.01-2μg/mL;

Immunohistochemistry: 5-40μg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


[IDENTIFICATION]




Western Blot; Sample: Porcine Skin
lysate Primary Ab: 0.2µg/ml Mouse Anti-
Human TPS Antibody Second Ab:
0.2µg/mL HRP-Linked Caprine Anti-
Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P; Sample:
Human Colorectal cancer Tissue;
Primary Ab: 40ug/ml Mouse Anti-
Human TPS Antibody Second Ab:
2µg/mL HRP-Linked Caprine Anti-
Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P;
Sample: Canine Small intestine Tissue;
Primary Ab: 10µg/ml Mouse Anti-
Human TPS Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P;
Sample: Canine Colon Tissue;
Primary Ab: 10µg/ml Mouse Anti-
Human TPS Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)



DAB staining on IHC-P;
Sample: Porcine Colon Tissue;
Primary Ab: 30ug/ml Mouse Anti-
Human TPS Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P;
Sample: Porcine Liver Tissue;
Primary Ab: 30ug/ml Mouse Anti-
Human TPS Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody
(Catalog: SAA544Mu19)




DAB staining on IHC-P;

Sample: Porcine Lung Tissue;
Primary Ab: 30ug/ml Mouse Anti-Human TPS Antibody
Second Ab: 2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)



DAB staining on IHC-P;

Sample: Porcine Lung Tissue
Primary Ab: 30µg/ml Mouse Anti-Human TPS Antibody
Control: Used PBS instead of primary antibody
Second Ab: 2µg/ml HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)



DAB staining on IHC-P;

Sample: Human Colorectal cancer Tissue
Primary Ab: 40µg/ml Mouse Anti-Human TPS Antibody
Control: Used PBS instead of primary antibody
Second Ab: 2µg/ml HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu19)

[**IMPORTANT NOTE**]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.