

MAA139Hu23

**Monoclonal Antibody to Tumor Necrosis Factor Related Apoptosis Inducing Ligand
(TRAIL)**

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

Purification: Protein A + Protein G affinity chromatography

Clone number: B5

Traits: Liquid

Concentration: 1mg/mL

UOM: 50µL

Cross Reactivity: Mouse;Rat

Applications: WB; ICC/IF

[IMMUNOGEN]

Immunogen: Recombinant TRAIL (Val114~Gly281) expressed in *E.coli*

Accession No.: RPA139Hu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunofluorescence: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no

obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]



Western Blot; Sample: Lane1: MCF7 cell lysate; Lane2: A549 cell lysate; Lane3: Rat Cerebrum lysate; Lane4: Mouse Cerebrum lysate Primary Ab: 0.05µg/ml Mouse Anti-Human TRAIL Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Mouse IgG Polyclonal Antibody (Catalog: SAA544Mu19)



AF488 staining on IF;

Sample: MCF7 cell
Primary Ab: 20µg/ml Mouse Anti-Human TRAIL Antibody
Second Ab: 2?g/ml AF488-Linked Caprine Anti-Mouse IgG Polyclonal Antibody
(Catalog: SAA544Mu11)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.