

PAH480Hu01

Polyclonal Antibody to 3-Oxoacid Coenzyme A Transferase 1 (OXCT1)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µL

Cross Reactivity: Mouse; Rat; Porcine.

Applications: WB; IHC.

[IMMUNOGEN]

Immunogen: Recombinant OXCT1 (Thr40~Leu489) expressed in E.coli

Accession No.: RPH480Hu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL

Immunohistochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Coud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]



Western Blot; Samples: Lane1: Human Western Blot; Sample: Lane1: Porcine Lung lysate; Lane2: Rat Cerebrum lysate; Lane3: Rat Heart lysate; Lane4: Rat Kidney lysate;


Primary Ab: 1µg/ml Rabbit Anti-Human **OXCT1** Antibody

Second Ab: 0.2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal **Antibody**


(Catalog: SAA544Rb19)


Heart lysate; Lane2: Porcine Kidney lysate; Lane3: Mouse Heart lysate; Lane4: Mouse Kidney lysate Primary Ab: 0.1µg/ml Rabbit Anti-**Human OXCT1 Antibody** Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal


> **Antibody** (Catalog: SAA544Rb19)




DAB staining on IHC-P; Samples: Human Liver Tissue; Primary Ab: 20µg/ml Rabbit Anti-Human OXCT1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)








DAB staining on IHC-P; Samples: Human Stomach Tissue; Primary Ab: 20µg/ml Rabbit Anti-Human OXCT1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Samples: Human Thyroid cancer Tissue; Primary Ab: 20µg/ml Rabbit Anti-Human OXCT1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal **Antibody** (Catalog: SAA544Rb19)

Coud-Clone Corp.




DAB staining on IHC-P; Samples: 20µg/ml Rabbit Anti-Human OXCT1 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P; Human Cerebrum Tissue; Primary Ab: Sample: Human Thyroid cancer Tissue Primary Ab: 20µg/ml Rabbit Anti-**Human OXCT1 Antibody** Control: Used PBS instead of primary antibody Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody

(Catalog: SAA544Rb19)



DAB staining on IHC-P; Sample: Human Kidney Tissue Primary Ab: 20µg/ml Rabbit Anti-Human OXCT1 Antibody Control: Used PBS instead of primary antibody Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P; Sample: Human Cerebrum Tissue Primary Ab: 20µg/ml Rabbit Anti-**Human OXCT1 Antibody** Control: Used PBS instead of primary antibody Second Ab: 2µg/ml HRP-Linked Caprine Anti-Rabbit IgG Polyclonal

Antibody (Catalog: SAA544Rb19)



[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.