

PAB120Rb51

Polyclonal Antibody to Arginase (ARG)

Organism Species: Oryctolagus cuniculus (Rabbit)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Guinea pig

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 200µL

Cross Reactivity: Human

Applications: IHC; ICC/IF

[IMMUNOGEN]

Immunogen: Recombinant ARG (Met1~Lys322) expressed in E.coli

Accession No.: RPB120Rb01

[APPLICATIONS]

Immunohistochemistry: 5-20µg/mL;

Immunofluorescence: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the



expiration date under appropriate storage condition.

[IDENTIFICATION]



DAB staining on IHC-P; Sample: Rabbit

Liver Tissue; Primary Ab: 20µg/ml Sample: Rabbit Kidney Tissue; AF488 staining on IF; Sample: HepG2 cell

Cavia Anti-Rabbit Arg Antibody Second Primary Ab: 20µg/ml Cavia Anti-Rabbit Primary Ab: 20µg/ml Cavia Anti-Rabbit Arg Antibody

DAB staining on IHC-P;

Ab: 2µg/mL HRP-Linked Rabbit Anti-

Cavia IgG Polyclonal Antibody (Catalog: SAA544Gu09)

Second Ab: 2µg/mL HRP-Linked Rabbit Anti-Cavia IgG Polyclonal Antibody

(Catalog: SAA544Gu09)

Arg Antibody

Second Ab: 2?g/ml AF488-Linked

Rabbit Anti-Cavia IgG Polyclonal

Antibody

(Catalog: SAA544Gu01)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.