

PAB231Ra01

Polyclonal Antibody to Cytokeratin 18 (CK18)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.31mg/mL

UOM: 20µL

Cross Reactivity: Human

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant CK18 (Glu72~Gly380) expressed in E.coli

Accession No.: RPB231Ra01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN3, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the


Cloud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]




DAB staining on IHC-P; Sample: Rat Stomach Tissue; Primary Ab: 20ug/ml Rabbit Anti-Rat CK18 Antibody Second Ab: 2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)




DAB staining on IHC-P;
Samples: Rat Small intestine Tissue;
Primary Ab: 30µg/ml Rabbit Anti-Rat
KRT18 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Rat Lung Tissue;
Primary Ab: 20?g/ml Rabbit Anti-Rat
CK18 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;

DAB staining on IHC-P;
Sample: Rat Kidney Tissue;
Primary Ab: 20?g/ml Rabbit Anti-Rat
CK18 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Sample: Rat Liver Tissue;
Primary Ab: 20?g/ml Rabbit Anti-Rat
CK18 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

DAB staining on IHC-P;
Sample: Rat Pancreas Tissue;
Primary Ab: 20?g/ml Rabbit Anti-Rat
CK18 Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

Coud-Clone Corp.




Figure. Western Blot; Sample: Lane1:
Human MCF-7 Cells; Lane2: Human
SGC7901 Cells; Lane3: Human Hela
Cells.

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.