

PAB239Hu06

Polyclonal Antibody to Cytokeratin 19 (CK19)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µL

Cross Reactivity: Porcine

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant CK19 (Asp244~Leu400) expressed in E.coli

Accession No.: RPB239Hu06

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the


Coud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]




Western Blot; Samples: Lane1: HepG2
cell lysate; Lane2: MCF7 cell lysate;
Lane3: Human Serum; Primary Ab:
0.1µg/ml Rabbit Anti-Human CK19
Antibody Second Ab: 0.2?g/ml HRPLinked Caprine Anti-Rabbit lgG
Polyclonal Antibody (Catalog:
SAA544Rb19)




DAB staining on IHC-P;
Sample: Porcine Uterus Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman CK19 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody (Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Small intestine Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman CK19 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody




(Catalog: SAA544Rb19)


Western Blot; Sample: Human Lung
lysate
Primary Ab: 0.01µg/ml Rabbit AntiHuman CK19 Antibody

Second Ab: 0.2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Sample: Porcine Pancreas Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman CK19 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody



DAB staining on IHC-P;

Sample: Porcine Liver Tissue
Primary Ab: 10µg/ml Rabbit AntiHuman CK19 Antibody
Control: Used PBS instead of primary
antibody
Second Ab: 2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody



(Catalog: SAA544Rb19) (Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.