

PAA764Mu01

Polyclonal Antibody to Forkhead Box Protein O1 (FOXO1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.46mg/mL

UOM: 100µL

Cross Reactivity: Human; Rat

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant FOXO1 (Arg222~Pro469) expressed in E.coli

Accession No.: RPA764Mu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Coud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]



DAB staining on IHC-P; Samples:

Mouse Skeletal muscle Tissue; Primary

Ab: 20?g/ml Rabbit Anti-Mouse FOXO1 Primary Ab: 20?g/ml Rabbit Anti-Mouse

Antibody Second Ab: 2µg/mL HRP-

Linked Caprine Anti-Rabbit IgG

Polyclonal Antibody (Catalog:

SAA544Rb19)

DAB staining on IHC-P;

Samples: Mouse Testis Tissue;


FOXO1 Antibody

Second Ab: 2µg/mL HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)



DAB staining on IHC-P;

Sample: Mouse Testis Tissue

Primary Ab: 20µg/ml Rabbit Anti-Mouse

FOXO1 Antibody

Control: Used PBS instead of primary


antibody

Second Ab: 2µg/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)





DAB staining on IHC-P;

Primary Ab: 20µg/ml Rabbit Anti-Mouse

FOXO1 Antibody


Control: Used PBS instead of primary

antibody

Second Ab: 2µg/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody



Western Blot; Samples: Lane1: Mouse

Sample: Mouse Skeletal muscle Tissue Liver lysate; Lane2: Rat Liver lysate;

Lane3: Hela cell lysate;

Primary Ab: 1µg/ml Rabbit Anti-Mouse

FOXO1 Antibody

Second Ab: 0.2?g/ml HRP-Linked

Caprine Anti-Rabbit IgG Polyclonal

Antibody

(Catalog: SAA544Rb19)



(Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.