

PA	Δ	RR	12	Лı	ı∩1
		UU		/I L	4 V I

Polyclonal Antibody to Peroxisome Proliferator Activated Receptor Gamma (PPARg)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µL

Cross Reactivity: Human; Rat

Applications: WB

[IMMUNOGEN]

Immunogen: Recombinant PPARg (Phe149~Ser273) expressed in E.coli

Accession No.: RPA886Mu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.



[IDENTIFICATION]



Western Blot; Sample: Rat Liver lysate; Western Blot; Sample: Rat Liver lysate; Cerebrum lysate; Lane2: K562 cell Primary Ab: 1µg/ml Rabbit Anti-Mouse Primary Ab: 1µg/ml Rabbit Anti-Mouse lysate; Primary Ab: 0.2µg/ml Rabbit Anti-PPARg Antibody **PPARg Antibody** Mouse PPARg Antibody Second Ab: Second Ab: 0.2µg/mL HRP-Linked Second Ab: 0.2µg/mL HRP-Linked 0.2?g/ml HRP-Linked Caprine Anti-Caprine Anti-Rabbit IgG Polyclonal Caprine Anti-Rabbit IgG Polyclonal Rabbit IgG Polyclonal Antibody Antibody Antibody (Catalog: SAA544Rb19) (Catalog: SAA544Rb19) (Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.