

PAA984Hu01

Polyclonal Antibody to Protein Tyrosine Phosphatase Receptor Type H (PTPRH)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 50µL

Cross Reactivity: Porcine

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant PTPRH (Asn844~Asp1096) expressed in E.coli

Accession No.: RPA984Hu01

[APPLICATIONS]

Western blotting: 0.01-3µg/mL;

Immunohistochemistry: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

Coud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]





DAB staining on IHC-P; Samples:
Human Liver Tissue; Primary Ab:
10μg/ml Rabbit Anti-Human PTPRH
Antibody Second Ab: 2μg/mL HRPLinked Caprine Anti-Rabbit IgG
Polyclonal Antibody (Catalog:

SAA544Rb19)

Western Blot; Sample: Lane1: Hela cell lysate; Lane2: Porcine Cerebrum lysate Primary Ab: 3µg/ml Rabbit Anti-Human PTPRH Antibody
Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody
(Catalog: SAA544Rb19)

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.