

PAB312Mu01

Polyclonal Antibody to Secretory Leukocyte Peptidase Inhibitor (SLPI)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 100µL

Cross Reactivity: Human

Applications: WB; IHC

[IMMUNOGEN]

Immunogen: Recombinant SLPI (Pro20~Met131) expressed in E.coli

Accession No.: RPB312Mu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-30µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the


Coud-Clone Corp.

expiration date under appropriate storage condition.

[IDENTIFICATION]




DAB staining on IHC-P; Samples:
Mouse Uterus Tissue; Primary Ab:
30µg/ml Rabbit Anti-Mouse SLPI
Antibody Second Ab: 2µg/mL HRPLinked Caprine Anti-Rabbit IgG
Polyclonal Antibody (Catalog:
SAA544Rb19)



DAB staining on IHC-P; DAB staining on IHC-P;
Samples: Mouse Esophagus Tissue; Samples: Mouse Testis Tissue;
Primary Ab: 30µg/ml Rabbit Anti-Mouse Primary Ab: 30µg/ml Rabbit Anti-Mouse






Samples: Mouse Testis Tissue;
Primary Ab: 30µg/ml Rabbit Anti-Mouse
SLPI Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



DAB staining on IHC-P;
Samples: Mouse Stomach Tissue;
Primary Ab: 30µg/ml Rabbit Anti-Mouse
SLPI Antibody
Second Ab: 2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Human Saliva
Primary Ab: 0.3µg/ml Rabbit AntiMouse SLPI Antibody
Second Ab: 0.2?g/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)

[IMPORTANT NOTE]



The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.