

PAA928Hu01

Polyclonal Antibody to Tumor Protein p53 (P53)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Polyclonal antibody preparation

Host: Rabbit

Purification: Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Traits: Liquid

Concentration: 0.5mg/mL

UOM: 1mL

Cross Reactivity: Rat;Porcine

Applications: WB; IHC; ICC/IF

[IMMUNOGEN]

Immunogen: Recombinant P53 (Gly108~Lys370) expressed in *E.coli*

Accession No.: RPA928Hu01

[APPLICATIONS]

Western blotting: 0.01-2µg/mL;

Immunohistochemistry: 5-20µg/mL;

Immunofluorescence: 5-20µg/mL;

Optimal working dilutions must be determined by end user.

[FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.


Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no


obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[IDENTIFICATION]



FITC staining on IF; Samples: Human


HepG2 cell; Primary Ab: 20μg/ml
Rabbit Anti-Human TP53 Antibody
Second Ab: 1.5μg/ml FITC-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody (Catalog: SAA544Rb18)



DAB staining on IHC-P;

Sample: Porcine Colon Tissue
Primary Ab: 10μg/ml Rabbit Anti-
Human P53 Antibody
Control: Used PBS instead of primary
antibody


Second Ab: 2μg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)




DAB staining on IHC-P;

Sample: Porcine Lymph node Tissue
Primary Ab: 10μg/ml Rabbit Anti-
Human P53 Antibody
Control: Used PBS instead of primary
antibody


Second Ab: 2μg/ml HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Lane1: Rat
Spleen lysate; Lane2: Human 293T cell
lysate
Primary Ab: 1μg/mL Rabbit Anti-Human
TP53 Antibody
Second Ab: 0.2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal



Knockout Varification:
Lane 1: Wild-type MCF7 cell lysate;
Lane 2: TP53 knockout MCF7 cell
lysate;
Predicted MW:
24,30,33,34,38,39,44kDa
Observed MW: 50kDa



Western Blot; Sample: Lane1: A431 cell
lysate; Lane2: MCF7 cell lysate
Primary Ab: 0.3μg/ml Rabbit Anti-
Human TP53 Antibody
Second Ab: 0.2μg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody


Antibody
(Catalog: SAA544Rb19)

Primary Ab: 1µg/ml Rabbit Anti-Human
TP53 Antibody

(Catalog: SAA544Rb19)

Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)



Western Blot; Sample: Porcine Skin
lysate

Primary Ab: 0.3µg/ml Rabbit Anti-
Human TP53 Antibody

Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody

(Catalog: SAA544Rb19)

[**IMPORTANT NOTE**]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.