

**PAD013Mu01**

**Polyclonal Antibody to X-Ray Repair Cross Complementing 5 (XRCC5)**

**Organism Species: *Mus musculus* (Mouse)**

***Instruction manual***

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

13th Edition (Revised in Aug, 2023)

### **[ PROPERTIES ]**

**Source:** Polyclonal antibody preparation

**Host:** Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity chromatography

**Traits:** Liquid

**Concentration:** 0.5mg/mL

**UOM:** 20μL

**Cross Reactivity:** Human

**Applications:** WB; ICC/IF

### **[ IMMUNOGEN ]**

**Immunogen:** Recombinant XRCC5 (Leu251~Ile732) expressed in *E.coli*

**Accession No.:** RPD013Mu01

### **[ APPLICATIONS ]**

Western blotting: 0.01-2μg/mL;

Immunofluorescence: 5-20μg/mL;

Optimal working dilutions must be determined by end user.

### **[ FORMULATION ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 24 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the

expiration date under appropriate storage condition.

**[ IDENTIFICATION ]**




Figure. FITC staining on IHC-P;  
Samples: Human HeLa Cells.




Figure. Western Blot; Sample: Lane1:  
Human K562 Cells; Lane2: Human  
HeLa Cells; Lane3: Human HepG2  
Cells.

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.