

**PAA473Ra01** 

Polyclonal Antibody to Xeroderma Pigmentosum, Complementation Group C (XPC)

Organism Species: Rattus norvegicus (Rat)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Polyclonal antibody preparation

Host: Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Traits: Liquid

Concentration: 0.5mg/mL

**UOM:** 200µL

Cross Reactivity: Human.

Applications: WB; FCM

#### [ IMMUNOGEN ]

Immunogen: Recombinant XPC (Glu734~Leu933) expressed in E.coli

Accession No.: RPA473Ra01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Flow cytometry: 20µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the


# Coud-Clone Corp.

expiration date under appropriate storage condition.

## [ IDENTIFICATION ]









Caprine Anti-Rabbit IgG Polyclonal

Antibody
(Catalog: SAA544Rb19)



Western Blot; Sample: Human Jurkat cell lysate;

Primary Ab: 1µg/ml Rabbit Anti-Rat
XPC Antibody
Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Rabbit IgG Polyclonal
Antibody
(Catalog: SAA544Rb19)



K562 human chronic myelogenous leukemia cell line was stained with rabbit Anti-rat XPC Polyclonal Antibody (Catalog # PAA473Ra01, filled red histogram) or Isotype control antibody (Catalog # IS067-Rb01, filled green histogram), followed by APC-conjugated Anti-rabbit IgG Secondary Antibody (Catalog # SAA544Rb15).

Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100.

## [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.