

RPB895Mu01 200µg

Recombinant Mannose Associated Serine Protease 1 (MASP1)

Organism Species: Mus musculus (Mouse)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: His25~Arg453

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 50µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 52.6kDa

Accurate Molecular Mass: 53kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-0.5 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]



		HTVELN	EMFGQIQSPG	YPDSYPSDSE
VTWNITVPEG	FRIKLYFMHF	NLESSYLCEY	DYVKVETEDQ	VLATFCGRET
TDTEQTPGQE	VVLSPGTFMS	VTFRSDFSNE	ERFTGFDAHY	MAVDVDECKE
REDEELSCOH	YCHNYIGGYY	CSCRFGYILH	TDNRTCRVEC	SGNLFTQRTG
TITSPDYPNP	YPKSSECSYT	IDLEEGFMVS	LQFEDIFDIE	DHPEVPCPYD
YIKIKAGSKV	WGPFCGEKSP	EPISTQTHSV	QILFRSDNSG	ENRGWRLSYR
AAGNECPKLQ	PPVYGKIEPS	QAVYSFKDQV	LVSCDTGYKV	LKDNEVMDTF
QIECLKDGAW	SNKIPTCKIV	DCGAPAGLKH	GLVTFSTRNN	LTTYKSEIRY
SCQQPYYKML	HNTTGVYTCS	AHGTWTNEVL	KRSLPTCLPV	CGVPKFSRKQ
ISR				

[IDENTIFICATION]




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.