

RPB319Hu01 200µg

Recombinant Microtubule Associated Protein 4 (MAP4)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



# [PROPERTIES]

**Source:** Prokaryotic expression

Host: E.coli

Residues: Met243~Val549

Tags: N-terminal His and GST Tag

**Subcellular Location:** Cytoplasm

**Purity:** > 80%

Traits: Freeze-dried powder

**Buffer formulation:** PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 4.5

Predicted Molecular Mass: 63.1kDa

**Accurate Molecular Mass:** 63kDa as determined by SDS-PAGE reducing conditions.

#### [USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

### [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ SEQUENCE ]

				MGLKTTDM
APSKETEMAL	AKDMALATKT	EVALAKDMES	PTKLDVTLAK	DMQPSMESDM
ALVKDMELPT	EKEVALVKDV	RWPTETDVSS	AKNVVLPTET	EVAPAKDVTL
LKETERASPI	KMDLAPSKDM	<b>GPPKENKKET</b>	ERASPIKMDL	APSKDMGPPK
ENKIVPAKDL	VLLSEIEVAQ	ANDIISSTEI	SSAEKVALSS	ETEVALARDM
TLPPETNVIL	TKDKALPLEA	EVAPVKDMAQ	LPETEIAPAK	DVAPSTVKEV
GLLKDMSPLS	ETEMALGKDV	<b>TPPPETEVVL</b>	IKNVCLPPEM	EVALTEDQV

## [ IDENTIFICATION ]




Figure . Gene Sequencing (extract)




Figure. SDS-PAGE

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.