

RPA684Hu02 10µg

Recombinant Mucin 5 Subtype B (MUC5B)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)

[PROPERTIES]

Source: Prokaryotic expression

Host: *E.coli*

Residues: Cys5653~Val5762

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 90%

Traits: Freeze-dried powder

Buffer formulation: 100mMNaHCO₃, 500mMNaCl, pH8.3, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 5.1

Predicted Molecular Mass: 15.7kDa

Accurate Molecular Mass: 16kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in ddH₂O to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

CEEDSCQV RINTTILWHQ GCETEVNITF CEGSCPGASK YSAEAQAMQH
QCTCCQERRV HEETVPLHCP NGSAILHTYT HVDECGCTPF CVPAPMAPPH
TRGFPAQEAT AV

[IDENTIFICATION]

TGTGAGGAGGCTCCTCTGTGAGTTCGATCAACAGACCACTCTGTGGCAGCAGGCTGTGGAGAGCCGAGGCTCAACATCACTCTCTGGAGGGCTCTGTGCGGAGAGCTGTCAGGTACTCAACAGAGGCGGAGGCGCATGCGAGCAACAGTGCACCTGTCTGCCAGGAGAGGCGGGTCCAGGAGGAGAG
 C E E D S C Q V R I N T T I L W H Q G C E T E V N I T F C E G S C P G A S K Y S A E A Q A M Q H Q C T C C Q E R R V H E E T




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.