

RPA709Hu01 200µg

Recombinant Toll Like Receptor 9 (TLR9)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



[PROPERTIES]

Source: Prokaryotic expression

Host: E.coli

Residues: Thr28~Ser362

Tags: N-terminal His Tag

Subcellular Location: Lysosome, Cytoplasm, Endoplasmic reticulum lumen, Endosome

Purity: > 80%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% Sarcosyl, 5% Trehalose.

Original Concentration: 450µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 9.1

Predicted Molecular Mass: 41.1kDa

Accurate Molecular Mass: 41kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
TLP AFLPCELQPH GLVNCNWLFL
KSVPHFSMAA PRGNVTSLSL SSNRIHHLHD SDFAHLPSLR HLNLKWNCPP
VGLSPMHFPC HMTIEPSTFL AVPTLEELNL SYNNIMTVPA LPKSLISLSL
SHTNILMLDS ASLAGLHALR FLFMDGNCYY KNPCRQALEV APGALLGLGN
LTHLSLKYNN LTVVPRNLPS SLEYLLLSYN RIVKLAPEDL ANLTALRVLD
VGGNCRRCDH APNPCMECPR HFPQLHPDTF SHLSRLEGLV LKDSSLSWLN
ASWFRGLGNL RVLDLSENFL YKCITKTKAF QGLTQLRKLN LSFNYQKRVS
FAHLSLAPSF GS
```

[IDENTIFICATION]




Figure. Gene Sequencing (Extract)




Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.