

APB090Bo01 100µg

Active Glutathione S Transferase Pi (GSTp)

Organism Species: *Bos taurus*; Bovine (Cattle)

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Met1~Gln210

Tags: N-terminal His-tag

Purity: >95%

Traits: Freeze-dried powder

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: PBS, pH7.4, containing 0.01% SKL, 5% Trehalose.

Original Concentration: 200µg/mL

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.4

Predicted Molecular Mass: 24.9kDa

Accurate Molecular Mass: 25kDa as determined by SDS-PAGE reducing conditions.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```
MPPYTIVYFP VQGRCEAMRM LLADQGQSWK EEVVAMQSWL QGPLKASCLY
GQLPKFQDGD LTLYQSNAIL RHLGRTLGLY GKDQQEAAALV DMVNDGVEDL
RCKYVSLIYT NYEAGKEDYV KALPQHLKPF ETLLSQNKGG QAFIVGDQIS
FADYNLLDLL RIHQVLAPSC LDSFPLLSAY VARLNSRPKL KAFLASPEHM
NRPINGNGKQ
```

[ACTIVITY]

GSTp (Glutathione S-transferase P) is an enzyme that plays an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. GSTP1 is identified as a CDK5 (Cyclin dependent kinase-5) regulatory protein, and is thought to regulates negatively CDK5 activity via p25/p35 translocation. GSTP1 catalyze the endogenous glutathione conjugation 1-Chloro-2,4-dinitrobenzene (CDNB), which can increase in the absorbance at 340 nm. The reaction was performed in adding 10 µl 200 mM glutathione (reduced) and 10 µl 100 mM CDNB in 980 µl 100 mM NaH₂PO₄ (pH7.0), rapidly mixed. Then add 50 µl mixed substrates to 50 µl different concentrations of recombinant bovine GSTP1, mix gently. Incubated at 37 °C for 5min, then read at a wavelength of 340 nm. The specific activity of recombinant bovine GSTP1 is 2000 pmol/min/µg.

Specific Activity (pmol/min/ug)=

Adjusted V_{max}^* (OD/min) x well volume (L) x 10^{12} pmol/mol

ext. coeff** ($M^{-1}cm^{-1}$) x path corr.*** (cm) x amount of enzyme (ug)

*Adjusted for Substrate Blank

**Using the extinction coefficient $9600 M^{-1}cm^{-1}$

***Using the path correction 1 cm

[IDENTIFICATION]




Figure 1. SDS-PAGE

Sample: Active recombinant GSTp, Cattle

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.