

APB575Hu01 100µg

Active Active Secondary Lymphoid Tissue Chemokine (SLC)

Organism Species: *Homo sapiens (Human)*

Instruction manual

FOR RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

1st Edition (Apr, 2016)

[PROPERTIES]

Source: Prokaryotic expression.

Host: *E. coli*

Residues: Ser24~Pro134

Tags: N-terminal His-tag

Purity: >98%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Buffer Formulation: 20mM Tris, 150mM NaCl, pH8.0, containing 0.05% sarcosyl and 5% trehalose.

Applications: Cell culture; Activity Assays.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 10.0

Predicted Molecular Mass: 13.5kDa

Accurate Molecular Mass: 18kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


[SEQUENCE]

SDGGAQD CCLKYSQRKI PAKVRSYRK
QEPSLGCSIP AILFLPRKRS QAELCADPKE LWVQLMQHL DKTPSPQKPA
QGCRKDRGAS KTGKKGKGSK GCKRTERSQT PKGP

[ACTIVITY]

Secondary Lymphoid-tissue Chemokine (SLC) is a recently identified CC chemokine that is constitutively expressed in various lymphoid tissues and is a potent and specific chemoattractant for lymphocytes. Thus, chemotaxis assay used 24-well microchemotaxis system was undertaken to detect the chemotactic effect of SLC on the human T-lymphocyte leukemia cell line Jurkat. Briefly, Jurkat cells were seeded into the upper chambers (100uL cell suspension, 10⁶ cells/mL in RPMI 1640 with FBS free) and SLC (1ng/mL, 15ng/mL and 150ng/mL diluted separately in serum free RPMI 1640) was added in lower chamber with a polycarbonate filter (8um pore size) used to separate the two compartments. After incubation at 37°C with 5% CO₂ for 1h, the filter was removed, then cells in low

chamber were observed by inverted microscope at low magnification ($\times 100$) and the number of migrated cells were counted at high magnification ($\times 400$) randomly (five fields for each filter). Result shows SLC is able to induce migration of Jurkat cells. The migrated Jurkat cells in low chamber at low magnification ($\times 100$) were shown in Figure 1. Five fields of each chamber were randomly chosen, and the migrated cells were counted at high magnification ($\times 400$). Statistical results were shown in Figure 2. The optimum chemotaxis of SLC occurs at 15-150ng/mL.



A

B

Figure 1. The chemotactic effect of SLC on Jurkat cells

(A) Jurkat cells were seeded into the upper chambers and serum free RPMI 1640 with 150ng/mL SLC was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 1h;

(B) Jurkat cells were seeded into the upper chambers and serum free RPMI 1640 without SLC was added in lower chamber, then cells in lower chamber were observed at low magnification ($\times 100$) after incubation for 1h.




Figure 2. The chemotactic effect of SLC on Jurkat cells.

[**IDENTIFICATION**]




Figure 3. SDS-PAGE

Sample: Active recombinant SLC, Human




Figure 4. Western Blot

Sample: Recombinant SLC, Human;

Antibody: Rabbit Anti-Human SLC Ab (PAB575Hu01)

[IMPORTANT NOTE]

The kit is designed for in vitro and research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.