



A90594Ra01 Polyclonal Antibody to Cytochrome C, Somatic (CYCS) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

[PRODUCT INFORMATION]

Immunogen: CYCS	Western Blot		
Clonality: Polyclonal	kDa 75		
Host: Rabbit	50		
Species Reactivity: Rat	37		
Ig type: Rabbit IgG	57		
Purification: Antibodies are purified by	25		
target protein affinity chromatography.	20		
Applications: WB, ICC, IHC-P, IHC-F	15		
Form: Liquid	10	-	←13kDa
Size: 100µg	10		

Sample: Recombinant rat CYCS

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant rat CYCS (Gln13~Leu99) expressed in *E.coli*. Molecular Weight: 11.1kDa USCN accession No.: P90594Ra01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-QKCAQCHT VEKGGKHKTG PNLHGLFGRK TGQAAGFSYT DANKNKGITW GEDTLMEYLE NPKKYIPGTK MIFAGIKKKG ERADLIAYL

Unique product Superb quality Client favorite Nicest service 💿 ISO9001:2008; 💿 ISO13485:2003; CE 11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R. China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com





[ANTIBODY SPEFICITY]

Anti CYCS is a rabbit polyclonal antibody raised against rat CYCS. This antibody has been selected for its ability to recognize rat CYCS in immunohistochemical staining and western blotting, non cross-reactive with other members of the family.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Optimal working dilutions must be determined by end user.

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH 7.4, containing 0.02%Na₃N, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant rat CYCS (Gln13~Leu99) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading buffer: 100mM Tris(PH 8.8), 2%SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, Na₃N 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

Unique product Superb quality Client favorite Nicest service (2) ISO9001:2008; (2) ISO13485:2003; (6)

11271 Richmond Avenue, H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Email: mail@uscnk.us | Http://www.uscnk.us Export Processing Zone, Wuhan 430056, P.R.China | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-9551 | Email: mail@uscnk.com | Http://www.uscnk.com