

PAA791Bo01

Polyclonal Antibody to Osteonectin (ON)
Organism Species: Bos taurus; Bovine (Cattle)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
 NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: Osteonectin, Bovine

Clonality: Polyclonal

Host: Rabbit


Immunoglobulin Type: IgG

Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg



Sample: Recombinant Osteonectin, Bovine

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant osteonectin (Ala18~Ile303) expressed in *E.coli*.

USCN Accession No.: RPA791Bo01

Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHSGS-APQ QEALPDETEV VEETVAEVAE VPVGANPVQV EVGEFDDGAE
 ETEEEVVAEN PCQNHCKHG KVCELDENNT PMCVCDPTS CPAIGEFK VCSNDNKTFD
 SSCHFFATKC TLEGTKKGHK LLDYIGPCK YIPPCLDSEL TEFPLMRDW LKNVLVTLYE
 RDEDNNLLTE KQKLRVKKIH ENEKRLEAGD HPVELLARDF EKNYNMYIFP VHWQFGQLDQ
 HPIDGYLSHT ELAPLRAPLI PMEHCCTTRFF ETCDLNDNDKY IALDEWAGCF GIKEKDIDKD LVI

[ANTIBODY SPECIFICITY]

The antibody is a rabbit polyclonal antibody raised against osteonectin. It has been selected for its ability to recognize osteonectin in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN_3 , 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant osteonectin (Ala18~Ile303) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemiluminescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN_3 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.