

PAA672Hu02**Polyclonal Antibody to Slit Homolog 2 (Slit2)****Organism Species: Homo sapiens (Human)*****Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PRODUCT INFORMATION]**Immunogen:** Slit2, Human**Clonality:** Polyclonal**Host:** Rabbit**Immunoglobulin Type:** IgG**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 200µg/mL**UOM:** 100µg**[IMMUNOGEN INFORMATION]****Immunogen:** Recombinant Slit2 (Cys1416~Val1528) with N-terminal His-Tag expressed in *E.coli*.**Accession No.:** RPA672Hu02**[ANTIBODY SPECIFICITY]**

The antibody is a rabbit polyclonal antibody raised against Slit2. It has been selected for its ability to recognize Slit2 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant Slit2 (Cys1416~Val1528) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.
5uL per well when used in enhanced chemilumescence (ECL).


Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.


[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

[IMAGES]



Used in Western Blot, Sample:
Recombinant Slit2, Human



Used in DAB staining on formalin fixed
paraffin-embedded liver cancer tissue

PAA672Hu02


Polyclonal Antibody to Slit Homolog 2 (Slit2)

Organism Species: Homo sapiens (Human)


Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY


NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES




**Used in DAB staining on formalin fixed
paraffin- embedded lung cancer tissue**



**Used in DAB staining on formalin fixed
paraffin- embedded skin cancer tissue**



**Used in DAB staining on formalin fixed paraffin-
embedded prostate glands cancer tissue**



**Used in DAB staining on formalin fixed
paraffin- embedded breast cancer tissue**

PAA672Hu02


Polyclonal Antibody to Slit Homolog 2 (Slit2)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



Used in DAB staining on formalin fixed paraffin-
embedded pancreas cancer tissue