#### PAA133Ca01 Polyclonal Antibody to Tumor Necrosis Factor Alpha (TNFa) Organism Species: Canis familiaris; Canine (Dog) Instruction manual

#### FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES


[ PRODUCT INFORMATION ] kDa 75 Immunogen: TNFa, Canine 50 **Clonality:** Polyclonal 37 Host: Rabbit 25 Immunoglobulin Type: IgG 20 **Purification:** Affinity Chromatography. 15 Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL 10 **UOM:** 100µg

# 9th Edition (Revised in Jul, 2013)

Sample: Recombinant TNFa, Canine

#### [IMMUNOGEN INFORMATION]

Immunogen: Recombinant TNFa (Leu63~Leu233) expressed in E.coli. USCN Accession No.: RPA133Ca01 Sequence: The target protein is fused with N-terminal His-Tag and its sequence is listed below. MGHHHHHHSGSEF-LPNGLQLI SPLAQTVKSS SRTPSDKPVA HVVANPEAEG QLQWLSRRAN ALLANGVELT DNQLIVPSDG LYLIYSQVLF KGQGCPSTHV LLTHTISRFA VSYQTKVNLL SAIKSPCQRE TPEGTEAKPW YEPIYLGGVF QLEKGDRLSA EINLPNYLDF **AESGQVYFGI IAL** 





### [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against TNFa. It has been selected for its ability to recognize TNFa in immunohistochemical staining and western blotting.

#### [APPLICATIONS]

Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

# [<u>CONTENTS</u>]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

### [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant TNFa (Leu63~Leu233) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

# [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.