#### PAK016Mu01 Polyclonal Antibody to Visual System Homeobox Protein 2 (VSX2) Organism Species: Mus musculus (Mouse) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

# [PRODUCT INFORMATION]

Immunogen: VSX2, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

### [IMMUNOGEN INFORMATION]

Immunogen: Recombinant VSX2 (Pro2~His135) expressed in E.coli.

USCN Accession No.: RPK016Mu01

**Sequence:** The target protein is fused with N-terminal His-Tag and its sequence is listed below.

MGHHHHHHSGSEF-PVDFNGYWK MLSNENFEEY LRALDVNVAL RKIANLLKPD KEIVQDGDHM IIRTLSTFRN YIMDFQVGKE FEEDLTGIDD RKCMTTVSWD GDKLQCVQKG EKEGRGWTQW IEGDELHLEM RAEGVICKQV FKKVH

# [ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against VSX2. It has been selected for its ability to recognize VSX2 in immunohistochemical staining and western blotting.


# [APPLICATIONS]


Western blotting: 1:100-400 Immunocytochemistry in formalin fixed cells: 1:100-500 Immunohistochemistry in formalin fixed frozen section: 1:100-500 Immunohistochemistry in paraffin section: 1:50-200 Enzyme-linked Immunosorbent Assay: 1:100-200

# Coud-Clone Corp.

Optimal working dilutions must be determined by end user.

### [<u>IMAGES</u>]





Used in Western Blot ,Sample: Recombinant VSX2,Mouse

#### Used in DAB staining on fromalin fixed

paraffin- embedded Liver tissue

## [CONTENTS]

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

# [QUALITY CONTROL]

**Content:** The quality control contains recombinant VSX2 (Pro2~His135) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$  0.02%.

# [<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.