

LAB062Hu72
Biotin-Linked Monoclonal Antibody
To Heat Shock 70kDa Protein 1 Like Protein (HSPA1L)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



### [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

**Purification:** Protein A/G Affinity Chromatography.

Clone number: 6-1#

Label: Biotin

Original Antibody: MAB062Hu22

Traits: Liquid

Concentration: 500µg/mL

**UOM**: 200µg

Applications: WB; ICC; IHC-P; IHC-F; IF; ELISA.

### [ IMMUNOGEN ]

Immunogen: Recombinant HSPA1L (Met1~Asp641) expressed in *E.coli*.

Accession No.: RPB062Hu01

# [APPLICATIONS]

Western blotting: 0.5-2ug/ml

Immunocytochemistry in formalin fixed cells: 5-20ug/ml

Immunohistochemistry in formalin fixed frozen section: 5-20ug/ml

Immunohistochemistry in paraffin section: 5-20ug/ml Enzyme-linked Immunosorbent Assay: 0.05-2ug/ml

Optimal working dilutions must be determined by end user.

### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

## [ QUALITY CONTROL ]

**Content:** The quality control contains recombinant HSPA1L disposed in loading buffer.



Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL).

**Note:** The quality control is specifically manufactured as the positive control.

Not used for other purposes.

Loading Buffer: 100mM Tris(pH6.8), 1% SDS, 150mM NaCl, 50% glycerol,

0.02% BPB, 50mM DTT and 0.02% NaN<sub>3</sub>.

## [STORAGE AND STABILITY]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.