

**RPA821Po01 10 $\mu$ g**  
**Recombinant C Reactive Protein (CRP)**  
**Organism Species: *Sus scrofa*; Porcine (Pig)**  
***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

## **[ PROPERTIES ]**

**Residues:** Gln20~Pro222

**Tags:** N-terminal His-Tag

**Accession:** O19062

**Host:** *E. coli*

**Subcellular Location:** Secreted.

**Purity:** >95%

**Endotoxin Level:** <1.0EU per 1 $\mu$ g

(determined by the LAL method).

**Formulation:** Supplied as lyophilized form in PBS,  
pH7.4, containing 5% sucrose, 0.01% sarcosyl.

**Predicted isoelectric point:** 6.2


**Predicted Molecular Mass:** 24.4kDa

**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

## **[ USAGE ]**

Reconstitute in sterile PBS, pH7.2-pH7.4.



## **[ STORAGE AND STABILITY ]**

**Storage: Avoid repeated freeze/thaw cycles.**

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

## **[ SEQUENCES ]**

The sequence of the target protein is listed below.

Q TDMIGKAFVF PKESENSYVS LTARLTKPLT AFTVCLRVYT DLNRDYSLFS YATKTQYNEI  
LLFRGKTAVY SISVGGADV VFKPHQSSEPM HFCMTWESTS GITELWVDGK PMVRRSLKRG  
YSLGTQASII LGQEQDAFAG GFEKNQCLVG DIGDVNMWDY VLSPEEINTV YAGGTFSPNV  
LNWRALRYEM SGEVYVKPQL WP

## **[ REFERENCES ]**

1. Chomdej S., *et al.* (2004) *Anim. Genet.* 35:469-470.
2. Pepys MB., Hirschfield GM. (2003) *J. Clin. Invest.* 111 (12): 1805–12.
3. Mantovani A., *et al.* (2008) *J. Clin. Immunol.* 28 (1): 1–13.
4. Zacho J., *et al.* (2008) *N. Engl. J. Med.* 359 (18): 1897–908.