

EPC534Mu61 100µg
Eukaryotic Histidine Rich Glycoprotein (HRG)
Organism Species: *Mus musculus* (Mouse)
Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)

[PROPERTIES]

Source: Eukaryotic expression

Host: 293F Cell

Residues: Leu19~Lys525

Tags: N-terminal His Tag

Subcellular Location: Secreted

Purity: > 97%

Traits: Freeze-dried powder

Buffer formulation: PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.

Original Concentration: 200µg/mL

Applications: Positive Control; Immunogen; SDS-PAGE; WB.

(May be suitable for use in other assays to be determined by the end user.)

Predicted isoelectric point: 7.3

Predicted Molecular Mass: 58.86kDa

Accurate Molecular Mass: 80/60/45/35kDa as determined by SDS-PAGE reducing conditions.

Phenomenon explanation:

The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.

[USAGE]

Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCE]

```

                LS PTNCDA SEPL AEKVL DLINK GRRSGYVFEL
LRVSDAHLDR AGTATVYYLA LDVIESDCW LSTKAQDDCL PSRWQSEIVI
GQCKVIATRY SNESQDLSVN GYNCTTSSVS SALRNTKDSP VLLDFFEDSE
LYRKQARKAL DKYKTDNGDF ASFRVERAER VIRARGGERT NYYVEFSMRN
CSTQHFRSP LVFGFCRALL SYSIETSDLE TPDSIDINCE VFNIEDHKDT
SDMKPHWGHE RPLCDKHLCK LSGSRDHHHT HKTDKLGCPP PPEGKDNSDR
PRLQEGALPQ LPPGYPPHSG ANRTHRPSYN HSCNEHPCHG HRPHGHHPHS
HHPGGHSHG HHPHGHHPHS HSHGHHPG HHPHGHHPHG HHPHGHHPHG
HHPHGDFLD YGPCDPPSNS QELKGQYHRG YGPPHGHSRK RGP GKGLFPF
HHQIGYVYR LPPLNIGEVL TLPEANFPSF SLPNCNRS LQ PEIQPFQTA
SRSCPGKFES EFPQISKFFG YTPPK
    
```

[IDENTIFICATION]




Figure. Gene Sequencing (Extract)



Figure. SDS-PAGE

[IMPORTANT NOTE]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.