

**PAN056Hu81**

**FITC-Linked Antibody**

**To T-Cell Immunoreceptor With Ig And ITIM Domains Protein (TIGIT)**

**Organism Species: Homo sapiens (Human)**

***Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

## **[ PRODUCT INFORMATION ]**

**Immunogen:** TIGIT, Human

**Clonality:** Polyclonal

**Conjugation:** FITC

**Host:** Rabbit


**Immunoglobulin Type:** IgG

**Purification:** Affinity Chromatography.

**Applications:** WB, ICC, IHC-P, IHC-F, ELISA

**Concentration:** 200µg/mL

**UOM:** 100µg



*Sample: Recombinant TIGIT, Human*

## **[ IMMUNOGEN INFORMATION ]**

**Immunogen:** Recombinant TIGIT (Gly21~Leu234) expressed in *E.coli*.

**Accession No.:** RPN056Hu01

**Sequence:** The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEF-GMMTGTIETT GNISAEKGGS  
IILQCHLSST TAQVTQVNWE QQDQLLAICN ADLGWHISPS FKDRVAPGPG LGLTLQSLTV  
NDTGEYFCIY HTYPDGTYTG RIFLEVLESS VAEHGARFQI PLLGAMAATL VVICTAVIVV  
VALTRKKKAL RIHSVEGDLR RKSAGQEEWS PSAPSPPGSC VQAEAAPAGL CGEQRGEDCA  
ELHDYFNVLS YRSL

## **[ ANTIBODY SPECIFICITY ]**

The antibody is a rabbit polyclonal antibody raised against TIGIT. It has been selected for its ability to recognize TIGIT in immunohistochemical staining and western blotting.

## **[ APPLICATIONS ]**

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100

Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

## **[ CONTENTS ]**

**Form & Buffer:** Supplied as solution form in PBS, pH7.4, containing 0.02% NaN<sub>3</sub>, 50% glycerol.

## **[ STORAGE ]**

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.

**Note:** *As fluorescence can photobleach when exposed to light, so the antibody must be protected from light.*