RPA399Hu01 10µg Recombinant High Mobility Group Protein 1 (HMG1) Organism Species: Homo sapiens (Human) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

C Cloud-Clone Corp.

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Met1~Glu215 kDa 70 Tags: N-terminal His-Tag 44 Accession: P09429 33 Host: E. coli 26 Subcellular Location: Nucleus, Chromosome, 22 **Purity:** >95% 18 Endotoxin Level: <1.0EU per 1µg 14 (determined by the LAL method). 10 **Formulation:** Supplied as lyophilized form in 20mM Tris, 15% SDS-PAGE 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% sarcosyl, 5% trehalose, and preservative. Predicted isoelectric point: 5.9 Predicted Molecular Mass: 26.4kDa Applications: SDS-PAGE; WB; ELISA; IP. (May be suitable for use in other assays to be determined by the end user.)

[<u>USAGE</u>]

Reconstitute in ddH₂O.

Coud-Clone Corp.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

MGKGDPKKPR GKMSSYAFFV QTCREEHKKK HPDASVNFSE FSKKCSERWK TMSAKEKGKF EDMAKADKAR YEREMKTYIP PKGETKKKFK DPNAPKRPPS AFFLFCSEYR PKIKGEHPGL SIGDVAKKLG EMWNNTAADD KQPYEKKAAK LKEKYEKDIA AYRAKGKPDA AKKGVVKAEK SKKKKEEEED EEDEEDEEEE EDEEDEDEEE DDDDE

[REFERENCES]

- 1. Wen L., et al. (1989) Nucleic Acids Res. 17:1197-1214.
- 2. Ferrari S., et al. (1996) Genomics 35:367-371.
- 3. Xiang Y.-Y., et al. (1997) Int. J. Cancer 74:1-6.
- 4. Kornblit B., et al. (2007) Tissue Antigens 70:151-156.