



# P90777Mu01 Inducible T-Cell Co Stimulator (ICOS) Organism: Mus musculus (Mouse) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

5th Edition (Revised in January, 2013)

kDa

94 66.2

45

33

26

20

15% SDS-PAGE

14.4

**Mouse ICOS** 

### [ DESCRIPTION ]

Protein Names: Inducible T-Cell Co Stimulator

**Synonyms:** ICOS, Ailim

Species: Mouse

Size: 100µg

Source: Escherichia coli-derived

**Subcellular Location:** Membrane; Single-pass type I

membrane protein.

# [PROPERTIES]

Residues: Asn35~Ile131 (Accession # Q9WVS0),

with N-terminal His-Tag.

Grade & Purity: >95%, 15kDa as determined by

SDS-PAGE reducing conditions.

Formulation: Supplied as lyophilized form in PBS, pH

7.4, containing 5% sucrose, 0.01% sarcosyl.

**Endotoxin Level:** <1.0 EU per  $1\mu g$  (determined by

the LAL method).

**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Predicted Molecular Mass: 12.5kDa

Predicted isoelectric point: 7.8





#### [PREPARATION]

Reconstitute in sterile PBS, pH7.2-pH7.4.

## [STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

#### [SEQUENCES]

The target protein is fused with N-terminal His-Tag, its sequence is listed below.

MGHHHHHHSGSEF- NGGVQI SCKYPETVQQ LKMRLFRERE VLCELTKTKG SGNAVSIKNP

MLCLYHLSNN SVSFFLNNPD SSQGSYYFCS LSIFDPPPFQ ERNLSGGYLH I