

RPA048Hu01 50µg Recombinant Inhibin Beta E (INHbE) Organism Species: Homo sapiens (Human) Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[PROPERTIES]

Residues: Thr237~Ser350
Tags: N-terminal His-Tag
Accession: P58166

Host: E. coli


Subcellular Location: Secreted.

Purity: >95%

Endotoxin Level: <1.0EU per 1μg (determined by the LAL method).

Formulation: Supplied as lyophilized form in 10mM


PBS, pH7.4, containing 1mM DTT, 5% trehalose,



Predicted Molecular Mass: 14.0kDa

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)



[USAGE]

Reconstitute in sterile ddH₂O.



[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

TPTC EPATPLCCRR DHYVDFQELG WRDWILQPEG YQLNYCSGQC PPHLAGSPGI AASFHSAVFS LLKANNPWPA STSCCVPTAR RPLSLLYLDH NGNVVKTDVP DMVVEACGCS

[REFERENCES]

- 1. Hashimoto O., et al. (2002) Mol. Cell. Endocrinol. 194:117-122.
- 2. Hashimoto O., et al. (2006) Biochem. Biophys. Res. Commun. 341:416-424.
- 3. Kimmich T., et al. (2010) Arch. Gynecol. Obstet. 282:185-191.
- 4. Bergauer F., et al. (2009) J. Mol. Histol. 40:353-359.