[USAGE]

Reconstitute in ddH₂O.

[PROPERTIES]

Residues: His20~Cys164

Tags: N-terminal His-Tag

Subcellular Location: Secreted.

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Accession: P49708

Host: E. coli

Purity: >95%

Predicted isoelectric point: 9.1

Predicted Molecular Mass: 18.2kDa

Applications: SDS-PAGE; WB; ELISA; IP.

0.01% sarcosyl, 5% trehalose, and preservative.

Formulation: Supplied as lyophilized form in 20mM Tris,

150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT,

(May be suitable for use in other assays to be determined by the end user.)

10th Edition (Revised in Jan, 2014)

kDa 94

66.2 45

33


26

20

14.4

15% SDS-PAGE

FOR IN VITRO USE AND RESEARCH USE ONLY



NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[<u>SEQUENCES</u>]

The sequence of the target protein is listed below.

H TASSLNLVQL QDDIDKLKAD FNSSHSDVAD GGPIIVEKLK NWTERNEKRI ILSQIVSMYL EMLENTDKSK PHIKHISEEL YTLKNNLPDG VKKVKDIMDL AKLPMNDLRI QRKAANELFS ILQKLVDPPS FKRKRSQSQR RCNC

[REFERENCES]

1. Kaiser P., et al. (1998) Gene 207:25-32.

- 2. Weining K.C., et al. (1996) Eur. J. Immunol. 26:2440-2447.
- 3. Digby M.R., Lowenthal J.W. (1995) J. Interferon Cytokine Res. 15:939-945.
- 4. Lv Y.Z., et al. (2001) Submitted to the EMBL/GenBank/DDBJ databases.