

CPA575Hu32 100µg
OVA Conjugated Motilin (MTL)
Organism Species: Homo sapiens (Human)
Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PROPERTIES]

Antigen: MTL-OVA

Residues: Synthetic Peptide

Predicted isoelectric point: 8.6

Predicted Molecular Mass: 1831.0Da

Purity: >95%

Endotoxin Level: <1.0EU per 1µg (determined by the LAL method).

Formulation: Supplied as lyophilized form in PBS.

Applications: SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

[RELEVANCE]

Motilin is secreted by endocrine M cells that are numerous in crypts of the small intestine, especially in the duodenum and jejunum. Motilin has 22 amino acids and molecular weight of 2698. The main function of motilin is to increase the migrating myoelectric complex component of gastrointestinal motility and stimulate the production of pepsin. Motilin is also called "housekeeper of the gut" because it improves peristalsis in the small intestine and clears out the gut to prepare for the next meal.

[USAGE]

Reconstitute in sterile ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The synthetic peptide's sequence is listed below.

GELQRMQEKERNKGQ