

RPD744Hu01 50µg**Recombinant Lipolysis Stimulated Lipoprotein Receptor (LSR)****Organism Species: Homo sapiens (Human)*****Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

10th Edition (Revised in Jan, 2014)

[**PROPERTIES**]

Residues: Met1~Gly259**Tags:** Two N-terminal Tags, His-tag and S-tag**Accession:** Q86X29**Host:** *E. coli***Subcellular Location:** Cell membrane;

Single-pass membrane protein.

Purity: >90%**Endotoxin Level:** <1.0EU per 1µg


(determined by the LAL method).

Formulation: Supplied as lyophilized form in 20mM Tris, 500mM NaCl, pH8.0, containing 1mM EDTA, 0.01% sarcosyl, 5% trehalose, and preservative.**Predicted isoelectric point:** 7.1**Predicted Molecular Mass:** 33.9kDa**Accurate Molecular Mass:** 23kDa as determined by SDS-PAGE reducing conditions.**Applications:** SDS-PAGE; WB; ELISA; IP.

(May be suitable for use in other assays to be determined by the end user.)

Note: The possible reasons that the actual band size differs from the predicted are as follows:

1. Splice variants: Alternative splicing may create different sized proteins from the same gene.
2. Relative charge: The composition of amino acids may affects the charge of the protein.
3. Post-translational modification: Phosphorylation, glycosylation, methylation etc.
4. Post-translation cleavage: Many proteins are synthesized as pro-proteins, and then cleaved to give the active form.
5. Polymerization of the target protein: Dimerization, multimerization etc.



[USAGE]

Reconstitute in ddH₂O.

[STORAGE AND STABILITY]

Storage: Avoid repeated freeze/thaw cycles.

Store at 2-8°C for one month.

Aliquot and store at -80°C for 12 months.

Stability Test: The thermal stability is described by the loss rate of the target protein. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. (Referring from China Biological Products Standard, which was calculated by the Arrhenius equation.) The loss of this protein is less than 5% within the expiration date under appropriate storage condition.

[SEQUENCES]

The sequence of the target protein is listed below.

MQQDGLGVGT RNGSGKGRSV HPSWPWCAPR PLRYFGRDAR ARRAQTAAMA
LLAGGLSRGL GSHPAAGRD AVFVWLLLS TWCTAPARAI QVTVSNPYHV VILFQPVTLP
CTYQMTSTPT QPIVIWKYKS FCRDRIADAF SPASVDNQLN AQLAAGNPGY NPYVECDQDSV
RTVRVATKQ GNAVTLGDYY QGRRITITGN ADLTFDQTAW GDSGVYYCSV VSAQDLQGNN
EAYAELIVLD WLFVVVVCLA AFLIFLLG

[REFERENCES]

1. Garcia J.M., *et al.* (2007) Clin. Cancer Res. 13:6351-6358.
2. Masuda S., *et al.* (2011) J. Cell. Sci. 124:548-555.
3. Papatheodorou P., *et al.* (2013) MBio 4:e00244-13.
4. Otsuki T., *et al.* (2005) DNA Res. 12:117-126.