

MAA573Hu21**Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)****Organism Species: Homo sapiens (Human)*****Instruction manual***

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]**Immunogen:** PIIINP-KLH**Clonality:** Monoclonal**Clone number:** C6**Host:** Mouse**Immunoglobulin Type:** IgG2b kappa**Purification:** Affinity Chromatography.**Applications:** WB, ICC, IHC-P, IHC-F, ELISA**Concentration:** 500µg/mL**UOM:** 200ug**[IMMUNOGEN INFORMATION]****Immunogen:** Synthetic Peptide, PIIINP conjugated to KLH.**Accession No.:** CPA573Hu31**[RELEVANCE]**

PIIINP is the amino terminal peptide of type III procollagen, released from the precursor peptide during the synthesis and deposition of type III collagen. There is evidence that serum PIIINP measurement is an effective non-invasive test for the detection and monitoring of methotrexate-induced liver fibrosis and cirrhosis, and serial measurements may reduce the need for liver biopsy. PIIINP has a molecular weight of 42 000 and contains three distinct domains: a triple-helical domain (Col 3) in the middle of the molecule, the Col 1-domain at the amino terminal and Col 2-domain at the carboxyterminal end of the propeptide.

[ANTIBODY SPECIFICITY]

The antibody is a mouse monoclonal antibody raised against PIIINP. It has been selected for its ability to recognize PIIINP in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:100-400

Immunocytochemistry in formalin fixed cells: 1:100-500

Immunohistochemistry in formalin fixed frozen section: 1:100-500

Immunohistochemistry in paraffin section: 1:50-200

Enzyme-linked Immunosorbent Assay: 1:100-5000

Optimal working dilutions must be determined by end user.


[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.


[STORAGE]

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.


[IMAGES]




Used in DAB staining on formalin fixed paraffin-embedded human brain tissue



Used in DAB staining on formalin fixed paraffin-embedded human prostate gland



Used in DAB staining on formalin fixed paraffin-embedded human prostate gland cancer tissue



Used in DAB staining on formalin fixed paraffin-embedded human breast cancer

MAA573Hu21


Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY

NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES



Used in DAB staining on formalin fixed
paraffin- embedded human kidney tissue