

MAC231Hu21

Monoclonal Antibody to Protein Kinase B Alpha (PKBa)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

13th Edition (Revised in Aug, 2023)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG1 Kappa

Purification: Antigen-specific affinity chromatography followed by Protein A affinity

chromatography

Clone number: C2

Traits: Liquid

Concentration: 1mg/mL

**UOM**: 100µL

**Cross Reactivity: Porcine** 

**Applications: WB** 

## [ IMMUNOGEN ]

Immunogen: Recombinant PKBa (Ser122~Thr443) expressed in E.coli

Accession No.: RPC231Hu01

## [ APPLICATIONS ]

Western blotting: 0.01-2µg/mL;

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

## [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 24 months.


**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no



obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]





Western Blot; Samples: Lane1: Hela
cell lysate; Lane2: MCF7 cell lysate;
Lane3: A549 cell lysate;
Primary Ab: 0.2µg/ml Mouse AntiHuman PKBa Antibody
Second Ab: 0.2µg/ml HRP-Linked

Antibody
(Catalog: SAA544Mu19)

Caprine Anti-Mouse IgG Polyclonal

Western Blot; Sample: Porcine
Cerebrum lysate
Primary Ab: 0.2µg/ml Mouse AntiHuman PKBa Antibody
Second Ab: 0.2µg/ml HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody

(Catalog: SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.