PAC400Mu01 Polyclonal Antibody to Carnitine Acetyltransferase (CRAT) Organism Species: Mus musculus (Mouse) *Instruction manual*

FOR IN VITRO USE AND RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: CRAT, Mouse Clonality: Polyclonal Host: Rabbit Immunoglobulin Type: IgG Purification: Affinity Chromatography. Applications: WB, ICC, IHC-P, IHC-F, ELISA Concentration: 200µg/mL UOM: 100µg

[IMMUNOGEN INFORMATION]

Immunogen: Recombinant CRAT (Met1~Asp430) expressed in E.coli.

Accession No.: RPC400Mu01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGS- MLAFAARTVV KPLGLLKPSS LMKVSGRFKA HQDALPRLPV PPLQQSLDYY LKALQPIVSE EEWAHTKQLV DEFQTSGGVG ERLQKGLERR AKKMENWLSE WWLKTAYLQF RQPVVIYSSP GVILPKQDFV DLQGQLRFAA KLIEGVLDFK SMIDNETLPV EFLGGQPLCM NQYYQILSSC RVPGPKQDSV VNFLKSKRPP THITVVHNYQ FFELDVYHSD GTPLTSDQIF VQLEKIWNSS LQSNKEPVGI LTSNHRNTWA KAYNNLIKDK VNRESVNSIQ KSIFTVCLDK QVPRVSDDVY RNHVAGQMLH GGGSKFNSGN RWFDKTLQFI VAEDGSCGMV YEHAAAEGPP IVALVDHVME YTKKPELVRS PMVPLPMPKK LRFNITPEIK NDIEKAKQNL SIMIQDLDIM MLTFHHFGKD FPKSEKLSPD

[ANTIBODY SPECIFITY]

The antibody is a rabbit polyclonal antibody raised against CRAT. It has been selected for its ability to recognize CRAT in immunohistochemical staining and western blotting.


[APPLICATIONS]


Designed by Cloud-Clone Corp., Assembled by Uscn Life Science Inc. ISO9001:2008; ISO13485:2003 11271 Richmond Avenue Suite H104, Houston, TX 77082, USA | Toll free: 001-888-960-7402 | Fax: 001-832-538-0088 | Http://www.cloud-clone.us | E-mail: mail@cloud-clone.us Export Processing Zone Building F, Wuhan, Hubel 430056, PRC | Toll free: 0086-800-880-0687 | Fax: 0086-27-8425-8551 | Http://www.uscnk.com | E-mail: mail@cloud-clone.us

Coud-Clone Corp.

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200 Optimal working dilutions must be determined by end user.

[<u>IMAGES</u>]





Used in DAB staining on fromalin fixed paraffin-embedded testis tissue

Used in Western Blot, Sample: Recombinant CRAT, Mouse

[<u>CONTENTS</u>]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant CRAT (Met1~Asp430) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate.

5uL per well when used in enhanced chemilumescent (ECL). **Note:** The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN $_3$ 0.02%.

[<u>STORAGE</u>]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.