

PAB823Rb51

Polyclonal Antibody to Complement Component 9 (C9)
Organism Species: Oryctolagus cuniculus (Rabbit)

Instruction manual

FOR IN VITRO USE AND RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

9th Edition (Revised in Jul, 2013)

[PRODUCT INFORMATION]

Immunogen: C9, Rabbit

Clonality: Polyclonal

Host: Cavia

Immunoglobulin Type: IgG


Purification: Affinity Chromatography.

Applications: WB, ICC, IHC-P, IHC-F, ELISA

Concentration: 200µg/mL

UOM: 100µg

[IMMUNOGEN INFORMATION]



Sample: Recombinant C9, Rabbit

Immunogen: Recombinant C9 (Pro139~Lys519) expressed in E.coli.

Accession No.: RPB823Rb01

Sequence: The target protein is fused with two N-terminal Tags, His-tag and

T7-tag and its sequence is listed below.

MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEFELRR Q- PR LTCRNREVQE SELARTAGYG INILGMDPLA TPFDNEYYHG LCDRVWDGNT LTHYRKPWNV AVLAYETKID KNFRTEYYEE QMQAFKSIIE EETSNFNANL ALKFTPTEAK ASKAEEASPK NKSLDDNDKG FSSKFQFSYS KNETYQLFLS YSSQKEKMFL LVKGIIQLGR FVMKNRGVML TNTFLDDIKS LPTTYEKGEY FAFLETYGTH YSSSGSLGGR YELIYVLDKA SMKEKGIELN DIKKCLGFDL DLSLNIPGKS AGLSLTGQAN KNNCLKSGHG NAVNITRANL IDDVISLIRG GTQKFAFELK



EKLLTKAKMV DVTDFINWAS SLSDAPVLIN QKLSPIYNLI PVKIKDAHQK RQNLERGIED YINEFSTKK

[ANTIBODY SPECIFITY]

The antibody is a cavia polyclonal antibody raised against C9. It has been selected for its ability to recognize C9 in immunohistochemical staining and western blotting.

[APPLICATIONS]

Western blotting: 1:50-400

Immunocytochemistry in formalin fixed cells: 1:50-500

Immunohistochemistry in formalin fixed frozen section: 1:50-500

Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-200

Optimal working dilutions must be determined by end user.

[CONTENTS]

Form & Buffer: Supplied as solution form in PBS, pH7.4, containing 0.02% NaN₃, 50% glycerol.

[QUALITY CONTROL]

Content: The quality control contains recombinant C9 (Pro139~Lys519) disposed in loading buffer.

Usage: 10uL per well when 3,3'-Diaminobenzidine(DAB) as the substrate. 5uL per well when used in enhanced chemilumescent (ECL).

Note: The quality control is specifically manufactured as the positive control. Not used for other purposes.

Loading Buffer: 100mM Tris(pH8.8), 2% SDS, 200mM NaCl, 50% glycerol, BPB 0.01%, NaN₃ 0.02%.

[STORAGE]

Store at 4°C for frequent use. Stored at -20°C to -80°C in a manual defrost freezer for one year without detectable loss of activity. Avoid repeated freeze-thaw cycles.